



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Deta

About Entrez

Display Abstract Sort Save Text Clip Add Order

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy

PubMed Services
Journal Browser
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

☐ 1: Endocrinology 1987 Aug;121(2):657-66

Related Articles, **Books**,
LinkOut

Structural and functional studies of the tryptic core of the human chorionic gonadotropin beta-subunit.

Birken S, Kolks MA, Amr S, Nisula B, Puett D.

The beta-subunit of hCG was digested with trypsin to produce a modified form of the subunit for structure-function and immunological studies. After digestion of hCG beta with trypsin, the residual disulfide-linked core was isolated and found to be lacking the carboxy-terminal peptide (residues 115-145) and to contain bond cleavages between residues 2-3, 43-44, 74-75, and 95-96. The locations of these bond cleavages within the disulfide-bridged core were identified by isolation of the following peptides after reduction and S-carboxymethylation of the trypsin beta-core: beta 1-43, beta 3-43, beta 44-74, beta 44-95, beta 75-95, and beta 96-114. The circular dichroic spectrum of the tryptic beta-core over the wavelength region of about 200-320 nm was similar to that of the native subunit. In addition, the tryptic beta-core retained nearly full immunopotency in both polyclonal and monoclonal competitive RIAs and could combine with complementary native alpha-subunit. The hybrid, composed of the tryptic beta-core and native alpha, was purified and displayed a molar potency of about 0.1% relative to intact hCG in both a radioreceptor assay and an adenylate cyclase assay. Thus, the hybrid retained little biological activity. Although the extensive bond cleavages in the tryptic beta-core did not appear to change its secondary and tertiary structure sufficiently to significantly alter the circular dichroic spectrum, the immunoreactivity, or the capability to combine with its alpha-subunit complement, the biological functional integrity of the tryptic beta-core-containing hybrid was essentially abolished. Hence, the tryptic beta-core provides a useful derivative for detailed structure-function studies aimed at defining the necessary determinants for subunit association, receptor binding, and subsequent biological actions.

PMID: 2439319 [PubMed - indexed for MEDLINE]

Display Abstract Sort Save Text Clip Add Order

[Write to the Help Desk](#)

NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

i686-pc-linux-gnu May 10 2002 10:49:42